Sheet 1 of 2 ATTORNEY DOCKET: 2026-4124

Serial No.: 08/231,565

SERIAL NO. U.S. DEPARTMENT OF COMMERCE ATTY. DOCKET NO. FORM PTO-1449 PATENT AND TRADEMARK OFFICE 08/231,565 2026-4124 INFORMATION DISCLOSURE CITATION APPLICANT(S) KAWAKAMI, et al. (Use several sheets if necessary) FILING DATE GROUP ART UNIT 1813 April 22, 1994 U.S. PATENT DOCUMENTS FILING DATE **EXAMINER** SUBCLASS IF APPROPRIATE CLASS DOCUMENT NUMBER DATE NAME INITIAL 08/31/94 Boon et al. 5342774 FOREIGN PATENT DOCUMENTS TRANSLATION DATE DOCUMENT NUMBER COUNTRY SUBCLASS YES NO CLASS 05/24/84 DE 33 41 367 A 1 Germany 08/23/95 0 66 83 50 A1 **EPO** 2 13 35 43 A 08/25/84 GB 08/24/95 **PCT** W0 95 22561 10/13/94 **PCT** WO 94 23067 WO 93 14189 07/22/93 **PCT** OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Papers, Etc.) Storkus, W.J. et al. (1993) "Identification of human melanoma peptides recognized by Class I SJH restricted tumor infiltrating T lymphocytes" J. Immunol. 151(7):3719-3727 Kawakami, Yutaka et al. (1994) "Cloning of the gene coding for a shared human melanoma antigen recognized by autologous T cells infiltrating into tumor" Proc. Natl. Acad. Sci. U.S.A. 91(9):3515-3519 Falk, K., et al.: "Allele-specific motifs revealed by sequencing of self-peptides eluted from MHC molecules." Nature 1991, 351:290-296.

Alluft 3/27/96

188021\_1

Sheet 2 of 2 ATTORNEY DOCKET: 2026-4124

Serial No.: 08/231,565

55H	Kubo, R., et al.: "Definition of specific peptide motifs for four major HLA-A Alleles." Jour Immunology 1994, 152:3913-3924.	rnal of
S5H	Parker, K., et al.: "Sequence motifs important for peptide binding to the human MHC class molecule. HLA-A2." 1992, J. Immunol:3580-3587.	l 
STH	Ruppert, J., et al.: "Prominent role of secondary anchor residues in peptide binding to HLA-molecules." Cell 1993, 74:929-937.	A2.1
504	Kawakami, Y., et al.: "Identification of human melanoma antigen recognized by trans-infiltrallymphocytes associated with in vivo tumor rejection." <i>Pro. Natl. Acad. Sci. USA</i> 1994, 91:66462.	
5574	Adema, G.J. et al., "Molecular characterization of the melanocyte lineage-specific antigen gp Journal of Biological Chemistry 1994, 269:20126-20133.	100."
55H	EMBL DATABASE ACCESSION NUMBER M32295: 26 11-90 Vogel A.: Human KD mela specific secreted glycoprotein MRNA 3'end'	anocyte
55H	Sette, et al., "Peptide Binding to the Most Frequent HLA-A Class I Alleles Measured by Quantitative Molecular Binding Assays" Molecular Immunology 31:813-822 (1994)	
5JH	Wölfel, et al., "Analysis of Antigens Recognized in Human Melanoma Cells by A2-Restricte Cytolytic T-Lymphocytes (CTL)" Int. J. Cancer 1993 55:237-244.	d
554	Wölfel, et al., "Isolation of Naturally Processed Peptides Recognized by Cytolytic Lymphocy (CTL) on Human Melanoma Cells In Association with HLA-A2.1" Int. J. Cancer 1994 413	
554	Slingluff, C.L., et al., "Recognition of Human Melanoma Cells by HLA-A2.1 Restricted Cy 7-Lymphocytes as Mediated by Six Strand Peptide Epitopes" <i>Journal of Immunology</i> 1993:15 2963	
5571	GENBANK ACCESSION NUMBER M77348 "Human PMEL 17 mRNA" January 8, 1995	
SOH	GENBANK ACCESSION NUMBER U06654 "Human Differentiation Antigen Melan-A Prof RNA" July 30, 1994	tein in
50H	GENBANK ACCESSION NUMBER S73003 "GP100 Melanocyte Lineage Specific Antigen/PMELL7 Honolog" January 25, 1995	
SOH	GENBANK ACCESSION NUMBER U01874 "Human ME20 in RNA" May 27, 1994	

EXAMINER DATE CONSIDERED 8/27/96

EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

				_									Sheet	1 0	۱ <u> </u>	
FORM PTO-1449 U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE								ATTY, DOCKET NO. 2026-4124 SERIAL NO. 08/231,565								
INFORMATION DISCLOSURE CITATION								APPLICANT	APPLICANT  Kawakami And Rosenberg							
(Use several sheets if necessary)							FILING DATE	FILING DATE April 22, 1994			GROUP To be Assigned					
							PAT	ENT	DOCUMENTS	<del></del>						
EXAMINER INITIAL		DOCUMENT NUMBER							DATE	DATE NAME			CLASS SUBCLASS IF APPRO			
STH		5 2		6	2	ı	7	7	11/16/93	Brown et al.						
										1						
					•	FC	REIG	N PAT	ENT DOCUM	ENTS						
		DOCUMENT NUMBER							DATE	E COUNTRY C			CLASS SUBCLASS		TRANSLATION	
+				<u> </u>	Γ_						+			YEŞ	NO	
			_	-		$\vdash$	+	$\vdash$		-	+				-	
			<u> </u>	_			+	ļ			+			-		
L			ОТН	ER DO	CUM	L_	S (Incl.	dina d	luthor Title De	ue, Pertinent Papers, 1					<u> </u>	
571		1	Cou	lie, P.	G. et	al. (19	993) "G			antigens recognized b		cytol	lytic T-lymphoc	ytes" J.	<del></del> ,	
SJH	-		Immunotherap. 14:104-109  Coulie P.G. et al. "A new gene coding for a differentiation antigen recognized by autologous cytolytic T lymphocytes on													
	-	5	HLA-A2 melanomas", J. Exp Med (1994) 180:35-42													
55H		·	Maresh, C.A. et al.: Cloning and expression of the gene for the melanoma associated ME20 antigen. DNA and Cell Biology, 1994; 13:87-95													
224		4.5	Cox, A.L., et al. "Identification of a peptide recognized by five melanoma-specific human cytotoxic T cell lines" Science 1994; 264:716-719.													
ट्यम			"Brichard, V., et al.: "The tyrosinase gene codes for an antigen recognized by autologous cytolytic T lymphocytes on HLA-A2 melanomas". J. Exp. Med. 1993; 178:489-495.													
20H		15.1	Gaugler, B., et al. "Human gene MAGE-3 codes for an antigen recognized on a melanoma by autologous cytolytic T lymphocytes". J. Exp. Med. 1994; 179:921-930.													
30H		T.J.	Traversari, C., et al.: "A nonapeptide encoded by human gene MAGE-1 is recognized on HLA-A1 by cytolytic T lymphocytes directed against umor antigen MZ2-E". J Exp. Med. 1992; 176:1453-1457.													
ટામ		FILE	Cellis, E., et al.: "Induction of anti-tumor cytotoxic T lymphocytes in normal humans using primary cultures and synthetic peptides epitopes". Proc. Natl. Acad. Sci. U.S.A. 1994; 91:2105-2109.													
5241		Th-	Boon, T.: "Toward a genetic analysis of tumor rejection antigens". Adv. Cancer Res. 1992; 58:177-210.													
201		117.	_							melanoma antigens. J						
20H		, h.		-cakker, A.B.H., et al.: Melanocyte lineage-specific antigen gp100 is recognized by melanocyte-derived tumor infiltrating lymphocytes. J. Exp. Med. 1994; 179:1005-1009.												
50.H		43	Wölfel, T., et al.: Two tyrosinase nonapeptides recognized on HLA-A2 melanomas by autologous cytolytic T. lymphocytes. Eur. J. Immunol. 1994; 24:759-764.													
55H		\;; <sub>a</sub>	Adema, G.J., et al.: Melanocyte lineage-specific antigens recognized by monoclonal antibodies NK1-beteb, HMB-50, and HMB-45 are encoded by a single cDNA. Am J. Pathol. 1993; 143:1579-1585.													
504	1	<u>.</u> Bys	<ol> <li>Kwon, B.S., et al.: A melanocyte-specific gene, Pmcl 17, maps near the silver coat color locus on mouse chromosome 10 and is in a syntenic region on human chromosome 12. Proc. Natl. Acad. Sci. USA 1991; 88:9228-9232.</li> </ol>													
55H			Rosenberg, S.A., et al.: Use of tumor infiltrating lymphocytes and interleukin-2 in the immunotherapy of patients with metastatic melanoma. Preliminary report. N. Engl. J. Med. 1988; 319:1676-1680.													
SIH	Ę	X	Kawakami, Y., et al.,: Shared human melanoma antigens. Recognition by tumor infiltrating lymphocytes in HLA-A2.1 transfected melanomas. <i>J Immunol</i> 1992; 148:638-643.													
5074		E.	&Van der Bruggen, et al.: A gene encoding an antigen recognized by cytolytic T. lymphocytes on a human melanoma.  Science 1991; 254:1643-1647.													
EXAMINER DATE CONSIDERED 8/24/94																
EXAMINER: Initiation of the considered. Include	tial if reference co de copy of this fo	nsider m wit	ed, wh	comm	or not unicat	citati ion to	on is in applica	confo	rmance with MF	EP 609; Draw line thr	ough cit	ation	if not in confor	mance and	I not	

FURM PTO-FB-AB20 (also form PTO-1449)